

## 3 DAYS BEFORE SURGERY

Drop		Breakfast	Mid-Day	Bedtime
<b>MOXIFLOXACIN</b> (Tan top)	1 Drop <u>3</u> times a day	✓	✓	✓
<b>PROLENSA/BROMSITE</b> (Grey top)	1 Drop <u>1</u> time a day	✓		

## WEEK 1: DAY 1 - DAY 7 AFTER SURGERY

Drop		Breakfast	Mid-Day	Bedtime
<b>MOXIFLOXACIN</b> (Tan top)	1 Drop <u>3</u> times a day	✓	✓	✓
<b>PREDNISOLONE</b> (Pink top)	1 Drop <u>3</u> times a day	✓	✓	✓
<b>PROLENSA/BROMSITE</b> (Grey top)	1 Drop <u>1</u> time a day	✓		

## WEEK 2: DAY 8 - DAY 14 AFTER SURGERY

Drop		Breakfast	Mid-Day	Bedtime
<b>PREDNISOLONE</b> (Pink top)	1 Drop <u>3</u> times a day	✓	✓	✓
<b>PROLENSA/BROMSITE</b> (Grey top)	1 Drop <u>1</u> time a day	✓		

## WEEK 3: DAY 15 - DAY 30 AFTER SURGERY

Drop		Breakfast	Mid-Day	Bedtime
<b>PREDNISOLONE</b> (Pink top)	1 Drop <u>2</u> times a day	✓		✓
<b>PROLENSA/BROMSITE</b> (Grey top)	1 Drop <u>1</u> time a day	✓		

**Please wait at least 2 minutes between drops.**